

Food and Drug Administration, HHS

§ 558.62

Finding Aids section of the printed volume and on GPO Access.

§ 558.59 Apramycin.

(a) *Approvals.* Type A articles to sponsors identified in § 510.600(c) of this chapter as follows:

(1) 000986 for 75 grams apramycin (as apramycin sulfate) per pound for use as in paragraph (d)(1) of this section.

(2) [Reserved]

(b) [Reserved]

(c) *Related tolerances.* See § 556.52 of this chapter.

(d) *Conditions of use*—(1) *Swine*—(i) *Amount.* 150 grams per ton.

(ii) *Indications for use.* For control of porcine colibacillosis (weanling pig scours) caused by susceptible strains of *Escherichia coli*.

(iii) *Limitations.* Use for 14 days. Withdraw 28 days before slaughter.

(2) [Reserved]

[51 FR 9190, Mar. 18, 1986]

§ 558.60 Arsanilate sodium.

(a) *Approvals.* Type A medicated articles: 20, 50, or 100 percent to 015565 in § 510.600(c) of this chapter.

(b) *Related tolerances.* See § 556.60 of this chapter.

(c) *Conditions of use.* (1) It is used for chickens and turkeys as follows:

(i) *Grams per ton.* 90 (0.01 percent).

(ii) *Indications for use.* For growth promotion and feed efficiency; improving pigmentation.

(iii) *Limitations.* Withdraw 5 days before slaughter; as sole source of organic arsenic.

(2) Arsanilate sodium may be used in accordance with the provisions of this section in the combinations provided as follows:

(i) Amprolium in accordance with § 558.55.

(ii) Zoalene in accordance with § 558.680.

[41 FR 10992, Mar. 15, 1976, as amended at 51 FR 7395, Mar. 3, 1986; 51 FR 33897, Sept. 24, 1986]

§ 558.62 Arsanilic acid.

(a) *Approvals.* Type A medicated articles to sponsors in § 510.600(c) of this chapter as follows:

(1) To 015565: 20, 50, and 100 percent for use as in the table in paragraph (c)(1), entry (ii), item 1; entry (ii), item 2; entry (iv); entry (vi); and entry (vii) of this section.

(2) To 015565: 20 percent for use as in paragraph (c)(1), entry (i); entry (ii), item 3 of this section.

(3) To 061133: 90 grams per pound arsanilic acid and 4.6 grams per pound erythromycin equivalents as erythromycin thiocyanate for use as in paragraph (c)(1), entry (iii); 90 grams per pound arsanilic acid and 9.25 grams per pound erythromycin equivalents as erythromycin thiocyanate for use as in paragraph (c)(1), entry (v).

(b) *Related tolerances.* See § 556.60 of this chapter.

(c) *Conditions of use.* (1) It is used as follows:

Arsanilic acid in grams per ton	Combination in grams per ton	Indications for use	Limitations	Sponsor
(i) 45 to 90	1. Growing chickens: For growth promotion and feed efficiency; improving pigmentation.	Withdraw 5 days before slaughter; as sole source of organic arsenic.	015565
	2. Growing turkeys: For growth promotion and feed efficiency; improving pigmentation.do	015565
	3. Growing swine: For increased rate of weight gain and improved feed efficiency.do	015565
(ii) 90	Swine: As an aid in control of swine dysentery (hemorrhagic enteritis, bloody dysentery).do	015565
(iii)	Erythromycin 4.6	Chickens: growth promotion and feed efficiency; improving pigmentation..	As erythromycin thiocyanate; withdraw 5 days before slaughter; as sole source of organic arsenic..	012487
(iv)	Erythromycin 4.6 to 18.5.	Chickens: growth promotion and feed efficiency; improving pigmentation.	As erythromycin thiocyanate; withdraw 5 days before slaughter; as sole source of organic arsenic.	15565